


5th International Conference of IGCP 588 and Fieldtrip Guide

SEISMIC AND NON-SEISMIC INFLUENCES ON COASTAL CHANGE IN ALASKA

May 3–10, 2014, South-Central Alaska, USA



SEISMIC AND NON-SEISMIC INFLUENCES ON COASTAL CHANGE IN ALASKA

**A Fieldtrip Guide and Conference Abstracts
5th International Conference of IGCP 588**

COMPILED BY:

Natasha Barlow

Sea Level Research Unit
Department of Geography
Durham University
South Road, Durham, DH1 3LE, UK
n.l.m.barlow@durham.ac.uk

Rich Koehler

Neotectonics Group
State of Alaska, Department of Natural Resources
Division of Geological & Geophysical Surveys
3354 College Road, Fairbanks, Alaska 99709-3707
907-451-5006
richard.koehler@alaska.gov
www.dggs.alaska.gov



*This guidebook was made possible with the publications support of
State of Alaska*

Department of Natural Resources
Division of Geological & Geophysical Surveys
3354 College Road, Fairbanks, AK 99709-3707
907-451-5000
dggspubs@alaska.gov
www.dggs.alaska.gov

COSEISMIC VS CLIMATIC EFFECTS IN THE RECORD OF RELATIVE SEA-LEVEL CHANGES: AN EXAMPLE FROM LAST INTERGLACIALS IN SE SPAIN

Bardají, T.¹, Cabero, A.², Lario, J.², Zazo, C.³, Silva, P.G.⁴, Goy, J.L.⁵, Dabrio, C.J.⁶

¹ U.D. Geología. Universidad de Alcalá. 28871-Alcalá de Henares (Madrid), España.

Corresponding author teresa.bardaji@uah.es

² Facultad de Ciencias, UNED. 28040-Madrid, España acabero@ccia.uned.es;

Javier.lario@ccia.uned.es

³ Dpto. Geología, Museo Nacional CC. Naturales, CSIC. 28006-Madrid, España.

mcnzc65@mncn.csic.es

⁴ Dpto. Geología, Universidad de Salamanca, EPTS de Avila. 05003-Ávila, España.

pgsilva@usal.es

⁵ Dpto. Geología, Facultad de Ciencias; Universidad de Salamanca. 37008 Salamanca, España. joselgoy@usal.es

⁶ Dpto. Estratigrafía, Facultad de CC. Geológicas, UCM. 28040-Madrid, España.

dabrio@ucm.es

Trying to decipher how palaeoseismic activity has influenced the altitudinal disposition of interglacial deposits is one of the main challenges that must be achieved in order to quantify relative sea-level changes between consecutive highstands. Likewise, identifying palaeoseismic features in an area with low instrumental seismic activity can implement the seismic record and contribute to better seismic hazard knowledge. In this sense, Cope basin (SE Spain) becomes a reference basin to undertake this kind of analysis, firstly because of excellent outcrop conditions and secondly because the sea-level record is undoubtedly conditioned here by both effects: climatically driven changes and local and regional seismotectonics.

This basin is located in the inner part of the tectonically active structure of the Aguilas Arc (Eastern Betics) and holds one of the most complete and better exposed sequences of Quaternary marine terraces in SE Spain. The chronology of the whole sequence has been achieved by means of palaeomagnetism (upper Matuyama–Brunhes transition), faunal content (*Strombus bubonius*, MIS7–MIS5 in Iberian Peninsula) and sedimentary facies (oolithic sediments, characteristic of MIS5e). Several faults active during the Quaternary cross the basin, promoting the development of structural blocks with differential uplift and, hence, different present-day altitude of marine terraces, particularly MIS7–MIS5 units. A detailed topographic leveling of the whole Quaternary sedimentary sequence carried out across the basin evidences that tectonics has been active both before and after the last Interglacial.

The studied sequence includes three different highstands: two littoral sedimentary units separated by terrestrial deposits, affected by liquefaction (palaeoseismic activity), and the whole sequence being later cut by a palaeocliff and a wave-cut platform. This wave-cut platform is related to a later lower sea level that could have been promoted either by a climatically driven lower sea level or by a coseismic uplift. The comparative altitudinal analysis of the three maximum transgressive heights allows discriminating between these two possibilities.

Acknowledgments: Spanish Projects: CGL2012-33430, CGL2012-37581-CO2-01, ACI-2009-1037 and UCM Res. Group 910198. It is a contribution of the Spanish Working Group of IGCP Project 588.